

## Screening for alcohol use disorder in high-prevalence demographics

Highmark Blue Cross Blue Shield of Western New York (Highmark BCBSWNY) partners with Amerigroup companies to administer certain services to Medicaid Managed Care (MMC) and Child Health Plus (CHPlus) members. Please note, this information is specific to the MMC and CHPlus programs only.

We encourage you to screen your patients for alcohol use disorder (AUD) who present in the high-prevalence demographics listed below.

### High-prevalence demographics

The lifetime prevalence of AUD in the U.S. population is approximately 29.1%. However, only 19.8% of people with AUD receive treatment. Prevalence of AUD is high in white and Indigenous people, younger men (age < 65), unmarried people, and those with low incomes.<sup>1</sup>

22.8 million people over the age of 12 reported having a substance use disorder (SUD) in 2019; AUD accounted for 63% of this population. An additional 12% presented with AUD and another SUD (excluding nicotine) according to the National Survey on Drug Use and Health (NSDUH).<sup>2</sup>

### AUD and COVID-19

Evidence suggests that alcohol consumption increased during the COVID-19 pandemic. One study found that 60% of respondents reported increased alcohol-intake.<sup>3</sup> In 2020, alcohol sales increased by 262% online and 21% in stores, which participants reported was due to increased stress, alcohol availability, and lockdown boredom.<sup>4</sup> This increase was most substantial between March to April 2020. The study suggests those most affected by COVID-19 (job loss, friend loss, family loss, and isolation) may be more at risk of AUD.<sup>3</sup>

### AUD co-occurring with mental health conditions

People with a variety of mental health conditions are at increased risk of developing an AUD or have an existing co-occurring AUD.<sup>5</sup> While the rates are higher for co-occurring disorders with mental health conditions, there is also a higher risk of greater severity and a worse prognosis for both the mental condition and AUD.

Trauma, including adverse childhood events (ACEs) and post-traumatic stress disorder (PTSD), are often precursors for AUD.<sup>6</sup> Traumatic brain injuries (TBIs) are also associated with AUD. Alcohol intoxication is one of the strongest predictors of a TBI. In addition, people with a TBI are more likely to abuse alcohol.<sup>7</sup>

In most co-occurring disorders, the mental health condition preceded the AUD. This indicates that people diagnosed with a mental health condition should be screened for AUD. Preventive work should begin at the onset of symptoms of a mental health condition.<sup>5</sup>

### <https://providerpublic.mybcbswny.com>

Amerigroup Partnership Plan, LLC provides management services for Highmark Blue Cross Blue Shield of Western New York's managed Medicaid. Amerigroup Partnership Plan, LLC brinda servicios administrativos para Medicaid administrado de Highmark Blue Cross Blue Shield of Western New York.

Highmark Blue Cross Blue Shield of Western New York is a trade name of Highmark Western and Northeastern New York Inc., an independent licensee of the Blue Cross Blue Shield Association. Highmark Blue Cross Blue Shield of Western New York es un nombre comercial de Highmark Western y Northeastern New York Inc., un licenciatario independiente de Blue Cross Blue Shield Association.

NYW-NL-0626-22 March 2022

## What if I need assistance?

If you have difficulty connecting patients with AUD to treatment, please contact Highmark BCBSWNY. If you have questions about this communication or need assistance with any other item, call Provider Services at **866-231-0847**.

1 Recovery Research Institute, 2019

<https://www.recoveryanswers.org/addiction-101/epidemiology>

2 Substance Abuse and Mental Health Services Administration, 2020

[Results from the 2019 National Survey on Drug Use and Health](#)

3 Pollard, Tucker, and Green (2020) <https://doi.org/10.1001/jamanetworkopen.2020.22942>

4 The Nielsen Company (2020) <https://www.nielsen.com/us/en/insights/article/2020/rebalancing-the-covid-19-effect-on-alcohol-sales>

5 Glantz et al., (2020) <https://doi.org/10.1016/j.addbeh.2019.106128>

6 Brady and Back (2012) <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3860395>

7 Weil, Corrigan, and Karelina (2018) <https://pubmed.ncbi.nlm.nih.gov/31198656>



**Email is the quickest and most direct way to receive important information from Highmark Blue Cross Blue Shield of Western New York.**

To start receiving email from us (including some sent in lieu of fax or mail), submit your information using the QR code to the right or via our online form (<https://bit.ly/39mJ3dc>).

