

# Clinical Criteria

**Subject:** Synagis (palivizumab) -NY Medicaid

**Document #:** ING-CC-0007-NY      **Publish Date:** 8/15/2021

**Status:** New      **Last Review Date:** 9/1/2021

## Table of Contents

[Overview](#)      [Coding](#)      [References](#)

[Clinical criteria](#)      [Document history](#)

## Overview

BlueCross BlueShield of Western New York partners with Amerigroup companies to administer certain services to Medicaid Managed Care (MMC) and Child Health Plus (CHPlus) members. Please note, this correspondence is specific to the MMC and CHPlus programs only.

This document addresses the use of Synagis (palivizumab), a monoclonal antibody approved by the Food and Drug Administration for prevention of serious lower respiratory tract disease caused by respiratory syncytial virus (RSV) infection in select pediatric individuals.

Randomized placebo-controlled clinical trials have demonstrated the safety and efficacy of Synagis in reducing hospitalizations due to RSV infection and in reductions in other measures of RSV infection severity for a very specific group of infants and children. Epidemiologic data indicate that the risk of severe RSV infection most likely to require hospitalization is greater in the presence of risk factors.

In 2014, the American Academy of Pediatrics (AAP) issued updated guidelines regarding the use of immune prophylaxis for RSV. AAP reaffirmed this guidance in 2019. A summary of the AAP RSV guidance is as follows:

<b>Preterm Infants without Chronic Lung Disease (CLD) of Prematurity or Congenital Heart Disease (CHD)</b>
<ul style="list-style-type: none"> <li>Infants born before 29 weeks, 0 days gestation in the first year of life</li> </ul>
<b>Preterm Infants with CLD</b>
<ul style="list-style-type: none"> <li>Infants born before 32 weeks, 0 days gestation and a requirement for &gt;21% oxygen for at least 28 days after birth in the first year of life</li> </ul>
<b>Infants with CHD</b>
<ul style="list-style-type: none"> <li>Prophylaxis may be administered in first year of life to certain infants with hemodynamically significant heart disease</li> <li>Consultation with a cardiologist if recommended for patients with cyanotic heart disease for prophylaxis decisions</li> </ul>
<b>Children with Anatomic Pulmonary Abnormalities or Neuromuscular Disorder</b>
<ul style="list-style-type: none"> <li>Prophylaxis may be considered in first year of life to children with pulmonary abnormality or neuromuscular disease that impairs the ability to clear secretions from the upper airways</li> </ul>
<b>Immunocompromised Children</b>
<ul style="list-style-type: none"> <li>Prophylaxis may be considered in children under 24 months who will be profoundly immunocompromised during the RSV season</li> </ul>

### <https://providerpublic.mycbswny.com>

Amerigroup Partnership Plan, LLC provides management services for BlueCross BlueShield of Western New York's managed Medicaid. Amerigroup Partnership Plan, LLC brinda servicios administrativos para Medicaid administrado de BlueCross BlueShield of Western New York. Highmark Blue Cross Blue Shield of Western New York and BlueCross BlueShield of Western New York are trade names of Highmark Western and Northeastern New York Inc., an independent licensee of the Blue Cross Blue Shield Association. Highmark Blue Cross Blue Shield of Western New York y BlueCross BlueShield of Western New York son nombres comerciales de Highmark Western y Northeastern New York Inc., un licenciario independiente de Blue Cross Blue Shield Association. NYWPEC-3170-21 October 2021

Children with Down Syndrome
<ul style="list-style-type: none"> <li>Insufficient data available to routinely recommend prophylaxis</li> </ul>
Children with Cystic Fibrosis
<ul style="list-style-type: none"> <li>Insufficient data available to routinely recommend prophylaxis</li> </ul>
Timing of Prophylaxis for Alaska Native and American Indian Infants
<ul style="list-style-type: none"> <li>Greater flexibility in use of prophylaxis as a result of potentially higher disease burden</li> <li>Use of government RSV surveillance data may be helpful in decision-making</li> </ul>
Discontinuation of Prophylaxis Among Children who Experience Breakthrough RSV Hospitalization
<ul style="list-style-type: none"> <li>Discontinue prophylaxis</li> </ul>
Prophylaxis in the Second Year of Life
<ul style="list-style-type: none"> <li>Recommended in children who require <math>\geq 28</math> days of supplemental oxygen after birth and continue to require medical intervention (supplemental oxygen, chronic corticosteroid therapy, diuretics)</li> </ul>
Number of Monthly Doses in Season
<ul style="list-style-type: none"> <li>Maximum of 5</li> </ul>
Other
<ul style="list-style-type: none"> <li>Prophylaxis is not recommended for prevention of primary asthma or reduction of subsequent wheezing episodes</li> <li>Prophylaxis is not recommended for prevention of nosocomial disease</li> <li>Not recommended for use in RSV treatment</li> </ul>

Because 5 monthly doses of Syngis will provide more than 6 months of adequate serum concentrations for most infants, administration should be limited to peak RSV seasons in the State of NY, of August 2021 to April 2022. Qualifying infants born during RSV season will need fewer than 5 doses for protection until the season ends.

Specific information about national and regional RSV trends, especially pertaining to the peak variations in Florida and Alaska, is available from the National Respiratory and Enteric Virus Surveillance System (NREVSS) at: <http://www.cdc.gov/surveillance/nrevss/rsv/index.html>.

## Clinical Criteria

When a drug is being reviewed for coverage under a member's medical benefit plan or is otherwise subject to clinical review (including prior authorization), the following criteria will be used to determine whether the drug meets any applicable medical necessity requirements for the intended/prescribed purpose.

### Synagis (palivizumab)

Requests for Synagis (palivizumab) may be approved if the following criteria are met (2014 AAP):

- I. A maximum of 5 doses of Synagis may be approved for **infants during the first RSV season within the first year of life** (infants in their first year of life who were administered RSV prophylaxis in April – September 2021 for coverage during a delayed RSV season may be evaluated under first RSV season criteria in 2021-22) with any of the following:
    - A. Born before 29 weeks, 0 days gestation (up to and including 28 weeks, 6 days) and younger than 12 months of age at the **start** of the RSV season; **OR**
    - B. Documentation is provided indicating chronic lung disease of prematurity defined as birth at less than 32 weeks, 0 days gestation and a requirement for greater than 21% oxygen for at least 28 days after birth (not including asthma, reactive airway disease and cystic fibrosis without significant symptoms); **OR**
    - C. Documentation is provided indicating hemodynamically significant congenital heart disease (including infants with acyanotic heart disease who are receiving medication to control congestive heart failure and will require cardiac surgical procedures and infants with moderate to severe pulmonary hypertension); **OR**
    - D. Documentation is provided indicating anatomic pulmonary abnormalities (for example, tracheal ring) or a neuromuscular condition that impairs the ability to clear secretions from the upper airway because of ineffective cough; **OR**
    - E. Documentation is provided indicating cystic fibrosis with clinical evidence of chronic lung disease or nutritional compromise (weight for length less than tenth percentile);
- OR**
- II. A maximum of 5 doses of Synagis may be approved for **children during their second RSV season** with any of the following:
    - A. Documentation is provided indicating the individual is a preterm infant born at less than 32 weeks, 0 days gestation who required at least 28 days of oxygen after birth AND continues to require medical intervention within 6 months of the start of the second RSV season (including supplemental oxygen, chronic corticosteroid therapy or diuretics); **OR**
    - B. Documentation is provided indicating cystic fibrosis with severe lung disease (history of hospitalization, abnormal chest x-ray or CT scan) or weight for length less than tenth percentile;

**OR**

- III. A maximum of 5 doses of Synagis may be approved for **children younger than 24 months of age** with any of the following:
- A. Documentation is provided indicating profound immunocompromised status (such as severe combined immunodeficiency, advanced acquired immunodeficiency syndrome, undergoing organ or hematopoietic stem cell transplant, or an absolute lymphocyte count of less than 100 cells/mm<sup>3</sup>); **OR**
  - B. Documentation is provided indicating the individual is undergoing cardiac transplantation;
- OR**
- IV. One additional dose of Synagis may be approved for children younger than 24 months of age who have approval for a course of treatment and who undergo cardiopulmonary bypass for a surgical procedure.

Synagis approval is limited to RSV season as determined by CDC surveillance data (<http://www.cdc.gov/surveillance/nrevss/rsv/index.html>) or local health department.

Synagis (palivizumab) may not be approved for any of the following:

- I. All other indications not included above; **OR**
- II. Continued RSV prophylaxis for children who experience breakthrough RSV hospitalization; **OR**
- III. Treatment of known RSV disease; **OR**
- IV. Children who reach 24 months of age prior to the beginning of RSV season; **OR**
- V. More than two seasons of RSV prophylaxis; **OR**
- VI. Primary asthma prevention or to reduce subsequent episodes of wheezing; **OR**
- VII. Children with surgically corrected congenital heart disease or hemodynamically insignificant heart disease (including secundum atrial septal defect, small ventricular septal defect, uncomplicated pulmonic stenosis, uncomplicated aortic stenosis, mild coarctation of the aorta, and patent ductus arteriosus) who do not otherwise meet approval criteria; **OR**
- VIII. Children with Down syndrome who do not otherwise meet approval criteria.

## Quantity Limits

### Synagis (palivizumab) Quantity Limit

Drug	Limit
Synagis (palivizumab) 50 mg, 100 mg vial	15 mg/kg once a month for up to 5 doses per RSV season
Override Criteria	
One additional dose may be approved for individuals undergoing cardiopulmonary bypass for a surgical procedure as noted in clinical criteria.	

## Coding

The following codes for treatments and procedures applicable to this document are included below for informational purposes. Inclusion or exclusion of a procedure, diagnosis or device code(s) does not constitute or imply member coverage or provider reimbursement policy. Please refer to the member's contract benefits in effect at the time of service to determine coverage or non-coverage of these services as it applies to an individual member.

### CPT

**90378** Respiratory syncytial virus, monoclonal antibody, recombinant, for intramuscular use, 50 mg, each [Synagis]

### HCPCS

**S9562** Home injectable therapy, palivizumab, including administrative services, professional pharmacy services, care coordination, and all necessary supplies and equipment (drugs and nursing visits coded separately), per diem

### ICD-10 Diagnosis

**P07.21-P07.26** Extreme prematurity  
**P07.31-P07.38** Preterm Newborn  
**P27.1** Bronchopulmonary dysplasia originating in the perinatal period  
**P27.8-P27.9** Other chronic respiratory diseases originating in the perinatal period  
**I42.9** Cardiomyopathy, unspecified  
**I50.9** Heart failure, unspecified  
**Q20.0-Q20.9** Congenital malformation of cardiac chambers and

	connections
<b>Q21.0-Q21.8</b>	Ventricular septal defects
<b>Q22.0</b>	Pulmonary valve atresia

## Document History

Revised: -9/1/2021

Document History:

- 9/1/2021 – New York specific policy created based on guidance released from NYSDOH.
- 8/20/2021 – Annual Review: Add note providing direction on how to address the delayed 2021 RSV season and to refer to CDC website for RSV season by region. Add exclusion for more than two seasons of prophylaxis. Wording and formatting changes. Coding reviewed: Added ICD-10-CM P07.21-P07.26, P07.31-P07.38, P27.1, P27.8-P27.9, I42.9, I50.9, Q20.0-Q20.9, Q21.0-Q21.8, Q22.0
- 11/30/2020 – Administrative update to add documentation requirements.
- 8/21/2020 – Annual Review: No changes. Coding Reviewed: No changes.
- 09/23/2019 – Administrative update to add drug specific quantity limit.
- 09/09/2019 - Annual Review: Wording and formatting changes. Coding reviewed: No changes.
- 08/17/2018 – Annual Review: Wording and formatting updates. Add in reference for criteria.

## References

1. American Academy of Pediatrics Committee on Infectious Diseases and Bronchiolitis Guidelines Committee. Updated guidance for palivizumab prophylaxis among infants and young children at increased risk of hospitalization for respiratory syncytial virus infection: Policy Statement. *Pediatrics*. 2014; 134(2):415-420. Erratum in: *Pediatrics*. 2014; 134(6):1221. Available at: <http://pediatrics.aappublications.org/content/134/2/415.full>. Accessed: July 8, 2021.
2. American Academy of Pediatrics Committee on Infectious Diseases and Bronchiolitis Guidelines Committee. Updated guidance for palivizumab prophylaxis among infants and young children at increased risk of hospitalization for respiratory syncytial virus infection: Technical Report. *Pediatrics*. 2014; 134(2):e620-e638. Available at: <http://pediatrics.aappublications.org/content/134/2/e620.full.pdf+html>. Accessed: July 8, 2021.
3. DailyMed. Package inserts. U.S. National Library of Medicine, National Institutes of Health website. <http://dailymed.nlm.nih.gov/dailymed/about.cfm>. Accessed: July 8, 2021.
4. DrugPoints® System [electronic version]. Truven Health Analytics, Greenwood Village, CO. Updated periodically.
5. Lexi-Comp ONLINE™ with AHFS™, Hudson, Ohio: Lexi-Comp, Inc.; 2021; Updated periodically.

Federal and state laws or requirements, contract language, and Plan utilization management programs or policies may take precedence over the application of this clinical criteria.

No part of this publication may be reproduced, stored in a retrieval system or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from the health plan.

© CPT Only – American Medical Association